

The codes used in the table CLINICAL, ordered by organ system

CLIN	VAR	DESC	CLIN	GROUP
AFE		Avascular necrosis in the femoral head		Bone
BFT		Bone fracture with assumed adequate trauma		
BFR		Bone fracture without adequate trauma		
BDE		DEXA scan (bone density) <i>fill-in FRAX</i>		
OST		Diagnosis of osteoporosis		
AOT		Other avascular necrosis		
END		Carotic endarterectomy		Cardio
CEH		Cerebral hemorrhage		
CEI		Cerebral infarction (stroke)		
ANG		Coronary angioplasty/stenting		
BYP		Coronary artery by-pass grafting		
DVT		Deep vein thrombosis		
HTR		Heart transplantation		
AMI		Myocardial infarction		
PRO		Procedures on other arteries		
PUL		Pulmonary embolism		
KBY		Kidney biopsy		Kidney
KTR		Kidney transplantation		
PDI		Permanent dialysis (expected to last > 1 month)		
BLE		Bleeding from gastric or esophageal varices		Liver
NAS		Diagnosis of NASH (non-alcoholic steatosis hepatitis)		
CIR		Diagnosis of cirrhosis (histology)		
PHY		Diagnosis of portal hypertension		
HEN		Hepatic encephalopathy (stage III or IV)		
HRE		Hepatorenal syndrome		
LTR		Liver transplantation		
PER		Spontaneous bacterial peritonis (with cirrhosis)		
MDE		DEXA scan (body composition)		Metabolisme
DIA		Diabetes mellitus, first diagnosis		
BFN		Bacterial Pneumonia		Other
PAN		Pancreatitis		